

ON IDENTITIES OF INFINITE DIMENSIONAL LIE SUPERALGEBRAS

DUŠAN REPOVŠ AND MIKHAIL ZAICEV

ABSTRACT. We study codimension growth of infinite dimensional Lie superalgebras over an algebraically closed field of characteristic zero. We prove that if a Lie superalgebra L is a Grassmann envelope of a finite dimensional simple Lie algebra then the PI-exponent of L exists and it is a positive integer.

1. INTRODUCTION

We shall consider algebras over a field F of characteristic zero. One of the approaches in the investigations of associative and non-associative algebras is to study numerical invariants associated with their identical relations. Given an algebra A , we can associate the sequence of its codimensions $\{c_n(A)\}_{n \in \mathbb{N}}$ (all notions and definitions will be given in the next section).

This sequence gives some information not only about identities of A but also about structure of A . For example, A is nilpotent if and only if $c_n(A) = 0$ for all large enough n . If A is an associative non-nilpotent F -algebra then A is commutative if and only if $c_n(A) = 1$ for all $n \geq 1$.

For an associative algebra A with a non-trivial polynomial identity the sequence $c_n(A)$ is exponentially bounded by the celebrated Regev's Theorem [20] while $c_n(A) = n!$ if A does not satisfy any non-trivial polynomial identity. In the non-associative case the sequence of codimensions may have even faster growth. For example, if A is an absolutely free algebra then

$$c_n(A) = a_n n!$$

where

$$a_n = \frac{1}{2} \binom{2n-2}{n-1}$$

is the Catalan number, i.e. the number of all possible arrangements of brackets in the word of length n .

For Lie algebra L the sequence $\{c_n(L)\}_{n \in \mathbb{N}}$ is not exponentially bounded in general even if L satisfies non-trivial Lie identities (see for example [18]). Nevertheless, a class of Lie algebras with exponentially bounded codimensions is sufficiently wide. It includes in particular, all finite dimensional algebras [1, 11], Kac-Moody algebras [23, 24], infinite dimensional simple Lie algebras of Cartan type [15], Virasoro algebra, and many others.

2010 *Mathematics Subject Classification.* Primary 17C05, 16P90; Secondary 16R10.

Key words and phrases. Polynomial identity, Lie algebra, codimensions, exponential growth.

The first author was supported by the Slovenian Research Agency grants P1-0292-0101 and J1-4144-0101. The second author was partially supported by RFBR grant No 13-01-00234a. We thank the referee for several comments and suggestions.

In the case when $\{c_n(A)\}_{n \in \mathbb{N}}$ is exponentially bounded, the upper and the lower limits of the sequence $\{\sqrt[n]{c_n(A)}\}_{n \in \mathbb{N}}$ exist and a natural question arises: does the ordinary limit

$$\lim_{n \rightarrow \infty} \sqrt[n]{c_n(A)}$$

exist? In case of existence we call this limit $\exp(A)$ or PI-exponent of A .

Amitsur conjectured in the 1980's that for any associative P.I. algebra such a limit exists and it is a non-negative integer. This conjecture was confirmed first for verbally prime P.I. algebras in [4, 21], and later in the general case in [8, 9]. For Lie algebras a series of positive results was obtained for finite dimensional algebras [6, 7, 25], for algebras with nilpotent commutator subalgebras [17], for affine Kac-Moody algebras [23, 24], and some other classes (see [16]). For Lie superalgebras there exist only partial results [26, 27, 30, 31].

On the other hand it was shown in [28] that there exists a Lie algebra L with

$$3.1 < \liminf_{n \rightarrow \infty} \sqrt[n]{c_n(L)} \leq \limsup_{n \rightarrow \infty} \sqrt[n]{c_n(L)} < 3.9 \quad .$$

This algebra L is soluble and almost nilpotent, i.e. it contains a nilpotent ideal of finite codimension. In the general non-associative case there exists, for any real number $\alpha > 1$, an algebra A_α such that

$$\lim_{n \rightarrow \infty} \sqrt[n]{c_n(A_\alpha)} = \alpha.$$

(see [5]). Note also that by a recent result [12] there exist finite dimensional Lie superalgebras with a fractional limit $\sqrt[n]{c_n(L)}$.

In the present paper we shall study Grassmann envelopes of finite dimensional simple Lie algebras. Our main result is the following theorem:

Theorem 1. *Let $L_0 \oplus L_1$ be a finite dimensional simple Lie algebra over an algebraically closed field F of characteristic zero with some \mathbb{Z}_2 -grading. Let also $\tilde{L} = L_0 \otimes G_0 \oplus L_1 \otimes G_1$ be the Grassmann envelope of L . Then the limit*

$$\exp(\tilde{L}) = \lim_{n \rightarrow \infty} \sqrt[n]{c_n(\tilde{L})}$$

exists and is a positive integer. Moreover, $\exp(\tilde{L}) = \dim L$.

Another result of our paper concerns graded identities. Since any Lie superalgebra L is \mathbb{Z}_2 -graded one can consider \mathbb{Z}_2 -graded identities of L and the corresponding graded codimensions $c_n^{gr}(L)$. We shall prove that graded codimensions have similar properties.

Theorem 2. *Let $L = L_0 \oplus L_1$ be a finite dimensional simple Lie algebra over an algebraically closed field F of characteristic zero with some \mathbb{Z}_2 -grading. Let also $\tilde{L} = L_0 \otimes G_0 \oplus L_1 \otimes G_1$ be a Grassmann envelope of L . Then the limit*

$$\exp^{gr}(\tilde{L}) = \lim_{n \rightarrow \infty} \sqrt[n]{c_n^{gr}(\tilde{L})}$$

exists and is a non-negative integer. Moreover, $\exp^{gr}(\tilde{L}) = \dim L$.

In other words, both PI-exponent $\exp(\tilde{L})$ and graded PI-exponent $\exp^{gr}(\tilde{L})$ exist, they are integers and they coincide. Note that for an arbitrary \mathbb{Z}_2 -graded algebra the growth of ordinary codimensions and graded codimensions may differ. For example, if $A = M_k(F) \otimes F\mathbb{Z}_2$ with the canonical \mathbb{Z}_2 -grading induced from group algebra $F\mathbb{Z}_2$, where $M_k(F)$ is full $k \times k$ matrix algebra, then $\exp(A) = k^2$ while $\exp^{gr}(A) = 2k^2$ (see [10] for details). In the Lie case one can take $L = L_0 \oplus L_1$ to be a two-dimensional metabelian algebra with $L_0 = \langle e \rangle, L_1 = \langle f \rangle$ and with only one non-trivial product $[e, f] = f$. Then $c_n(L) = n - 1$ for all $n \geq 2$ hence $\exp(L) = 1$. On the other hand $\exp^{gr}(L) = 2$.

2. THE MAIN CONSTRUCTIONS AND DEFINITIONS

Let A be an arbitrary non-associative algebra over a field F and let $F\{X\}$ be an absolutely free F -algebra with a countable generating set X . A polynomial $f = f(x_1, \dots, x_n)$ is said to be an identity of A if $f(a_1, \dots, a_n) = 0$ for any $a_1, \dots, a_n \in A$. The set of all identities of L forms a T-ideal $Id(A)$ in $F\{X\}$, that is an ideal which is stable under all endomorphisms of $F\{X\}$. Denote by $P_n = P_n(x_1, \dots, x_n)$ the subspace of all multilinear polynomials on x_1, \dots, x_n in $F\{X\}$. Then $P_n \cap Id(A)$ is a subspace of all multilinear identities of A of degree n . In the case when $\text{char } F = 0$, the T-ideal $Id(A)$ is completely determined by the subspaces $\{P_n \cap Id(A)\}, n = 1, 2, \dots$.

For estimating how many identities an algebra A can have one can define the so-called n -th codimension of the identities of A or, for shortness, codimension of A :

$$c_n(A) = \dim \frac{P_n}{P_n \cap Id(A)}, \quad n = 1, 2, \dots$$

As it was mentioned above, the class of associative and non-associative algebras with exponentially bounded sequence $\{c_n(A)\}$ is sufficiently wide. In the case when $c_n(A) < a^n$ for some real a , one can define the lower and the upper PI-exponents of A as follows:

$$\underline{\exp}(A) = \liminf_{n \rightarrow \infty} \sqrt[n]{c_n(A)}, \quad \overline{\exp}(A) = \limsup_{n \rightarrow \infty} \sqrt[n]{c_n(A)}$$

and the ordinary PI-exponent

$$(1) \quad \exp(A) = \lim_{n \rightarrow \infty} \sqrt[n]{c_n(A)},$$

provided that $\underline{\exp}(A) = \overline{\exp}(A)$.

For \mathbb{Z}_2 -graded algebras one can also consider graded identities. Let X and Y be two infinite sets of variables and let $F\{X \cup Y\}$ be an absolutely free algebra generated by $X \cup Y$. If we suppose that all elements of X are even and all elements of Y are odd, i.e. $\deg(x) = 0, \deg(y) = 1$ for any $x \in X, y \in Y$ then $F\{X \cup Y\}$ can be naturally endowed by a \mathbb{Z}_2 -grading. A polynomial $f = f(x_1, \dots, x_m, y_1, \dots, y_n) \in F\{X \cup Y\}$ is said to be a graded identity of a superalgebra $A = A_0 \oplus A_1$ if $f(a_1, \dots, a_m, b_1, \dots, b_n) = 0$ for all $a_1, \dots, a_m \in A_0, b_1, \dots, b_n \in A_1$. Fix $0 \leq k \leq n$ and denote by $P_{k, n-k}$ the subspace of $F\{X \cup Y\}$ spanned by all multilinear polynomials in $x_1, \dots, x_k \in X, y_1, \dots, y_{n-k} \in Y$. Then $P_{k, n-k} \cap Id(A)$ is the set of all multilinear polynomial identities of the superalgebra $A = A_0 \oplus A_1$ in k even and $n - k$ odd variables.

One of the equivalent definitions of graded codimensions of A is

$$c_n^{gr}(A) = \sum_{k=0}^n \binom{n}{k} c_{k,n-k}(A),$$

where

$$c_{k,n-k}(A) = \dim \frac{P_{k,n-k}}{P_{k,n-k} \cap Id(A)}.$$

Starting from a \mathbb{Z}_2 -graded algebra of some class (Lie, Jordan alternative, etc.) one can construct a \mathbb{Z}_2 -graded algebra of different class using the notion of the Grassmann envelope. Grassmann envelopes play an exceptional role in PI-theory. For example, any variety of associative algebras is generated by the Grassmann envelope of some finite dimensional associative superalgebra [14]. In Lie case any so-called special variety is generated by the Grassmann envelope of a finitely generated Lie superalgebra [22].

We recall this construction for Lie and super Lie cases. Let G be the Grassmann algebra generated by 1 and the infinite set $\{e_1, e_2, \dots\}$ satisfying the following relations: $e_i e_j = -e_j e_i$, $i, j = 1, 2, \dots$. It is known that G has a natural \mathbb{Z}_2 -grading $G = G_0 \oplus G_1$ where

$$G_0 = \text{Span} \langle e_{i_1} \cdots e_{i_n} | n = 2k, k = 0, 1, \dots \rangle,$$

$$G_1 = \text{Span} \langle e_{i_1} \cdots e_{i_n} | n = 2k + 1, k = 0, 1, \dots \rangle.$$

Given a Lie algebra L with \mathbb{Z}_2 -grading $L = L_0 \oplus L_1$, its Grassmann envelope

$$G(L) = L_0 \otimes G_0 \oplus L_1 \otimes G_1 \subset L \otimes G$$

is a Lie superalgebra. Vice versa, if $L = L_0 \oplus L_1$ is a Lie superalgebra then $G(L)$ is an ordinary Lie algebra with a \mathbb{Z}_2 -grading.

3. COCHARACTERS OF GRASSMANN ENVELOPES

The main tool in studying codimensions asymptotics is representation theory of symmetric groups. We refer the reader to [13] for details. Symmetric group S_n acts naturally on multilinear polynomials in $F\{X\}$ as

$$(2) \quad \sigma f(x_1, \dots, x_n) = f(x_{\sigma(1)}, \dots, x_{\sigma(n)}).$$

Hence P_n is an FS_n -module and $P_n \cap Id(L)$ and also

$$P_n(L) = \frac{P_n}{P_n \cap Id(L)}$$

are FS_n -modules. S_n -character $\chi(P_n(L))$ is called n -th cocharacter of L and we shall write

$$\chi_n(L) = \chi(P_n(L)).$$

Recall that any irreducible FS_n -module corresponds to a partition λ of n , $\lambda \vdash n$, $\lambda = (\lambda_1, \dots, \lambda_k)$, where $\lambda_1 \geq \dots \geq \lambda_k$ are positive integers and $\lambda_1 + \dots + \lambda_k = n$. By the Maschke Theorem, any finite dimensional FS_n -module M decomposes into the direct sum of irreducible components and hence its character $\chi(M)$ has a decomposition

$$\chi(M) = \sum_{\lambda \vdash n} m_\lambda \chi_\lambda$$

where m_λ are non-negative integers. In particular, for the algebra L we have

$$(3) \quad \chi(L) = \sum_{\lambda \vdash n} m_\lambda \chi_\lambda.$$

Integers m_λ in (3) are called multiplicities of χ_λ in $\chi_n(L)$ and $d_\lambda = \deg \chi_\lambda = \chi_\lambda(1)$ are the dimensions of corresponding irreducible representations. Therefore

$$(4) \quad c_n(L) = \dim P_n(L) = \sum_{\lambda \vdash n} m_\lambda d_\lambda.$$

For any partition $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$ one can construct Young diagram D_λ containing λ_1 boxes in the first row, λ_2 boxes in the second row and so on:

$$D_\lambda = \begin{array}{|c|c|c|c|c|c|} \hline & & \cdots & & \cdots & \\ \hline & & \cdots & & & \\ \hline \vdots & & & & & \\ \hline & & & & & \\ \hline \end{array}$$

Given integers $k, l, d \geq 0$, we define the partition

$$h(k, l, d) = (\underbrace{l + d, \dots, l + d}_k, \underbrace{l, \dots, l}_d)$$

of $n = kl + d(k + l)$. The Young diagram associated with $h(k, l, d)$ is hook shaped, and we define $H(k, l)$, an infinite hook, as the union of all D_λ with $\lambda = h(k, l, d)$, $d = 1, 2, \dots$. For shortness we will say that a partition $\lambda \vdash n$ lies in the hook $H(k, l)$, $\lambda \in H(k, l)$, if $D_\lambda \subset H(k, l)$. In other words, $\lambda \in H(k, l)$ if $\lambda = (\lambda_1, \dots, \lambda_t)$ and $\lambda_{k+1} \leq l$. According to this definition we will say that the cocharacter of L lies in the hook $H(k, l)$ if $m_\lambda = 0$ in (3) as soon as $\lambda \notin H(k, l)$.

A particular case of $H(k, l)$ is an infinite strip $H(k, 0)$. In this case $\lambda \in H(k, 0)$ if $\lambda_{k+1} = 0$.

The following fact is well-known and we state it without proof.

Lemma 1. *Let L be a finite dimensional algebra, $\dim L = d < \infty$. Then $\chi_n(L)$ lies in the hook $H(d, 0)$ for all $n \geq 1$.*

□

Another important numerical invariant of the identities of L is the colength $l_n(L)$. By definition

$$(5) \quad l_n(L) = \sum_{\lambda \vdash n} m_\lambda$$

where m_λ are taken from (3). It easily follows from (4) and (5) that

$$(6) \quad \max\{d_\lambda | m_\lambda \neq 0\} \leq c_n(L) \leq l_n(L) \cdot \max\{d_\lambda | m_\lambda \neq 0\}.$$

For studying graded identities of $L = L_0 \oplus L_1$ we need to act separately on even and odd variables. More precisely, the space $P_{k,n-k} = P_{k,n-k}(x_1, \dots, x_k, y_1, \dots, y_{n-k})$ is an $S_k \times S_{n-k}$ -module where symmetric groups S_k, S_{n-k} act on x_1, \dots, x_k and y_1, \dots, y_{n-k} , respectively. Any irreducible $S_k \times S_{n-k}$ -module is a tensor product of S_k -module and an S_{n-k} -module and corresponds to the pair λ, μ of partitions, $\lambda \vdash k, \mu \vdash n - k$. As before, the subspace $P_{n-k} \cap Id(L)$ is an $S_k \times S_{n-k}$ -stable subspace and one can consider the quotient

$$P_{k,n-k}(L) = \frac{P_{k,n-k}}{P_{k,n-k} \cap Id(L)}$$

as an $S_k \times S_{n-k}$ -module. Its $S_k \times S_{n-k}$ -character $\chi_{k,n-k}(L) = \chi(P_{k,n-k}(L))$ is decomposed into irreducible components.

$$(7) \quad \chi_{k,n-k}(L) = \sum_{\substack{\lambda \vdash k \\ \mu \vdash n-k}} m_{\lambda,\mu} \chi_{\lambda,\mu}$$

and we define the $(k, n-k)$ -colength of L as

$$l_{k,n-k}(L) = \sum_{\substack{\lambda \vdash k \\ \mu \vdash n-k}} m_{\lambda,\mu}$$

with $m_{\lambda,\mu}$ taken from (7).

First, we prove some relations between graded and non-graded numerical invariants. We begin by recalling the correspondence between multilinear homogeneous polynomials in a free \mathbb{Z}_2 -graded Lie algebra and in a free Lie superalgebra. Let $f = f(x_1, \dots, x_k, y_1, \dots, y_m)$ be a non-associative polynomial multilinear on $x_1, \dots, x_k, y_1, \dots, y_m$, where x_1, \dots, x_k are supposed to be even and y_1, \dots, y_m odd indeterminates. Then f is a linear combination of monomials from $P_{k,m}$. Let $M = M(x_1, \dots, x_k, y_1, \dots, y_m)$ be such a monomial. We fix positions of y_1, \dots, y_m in M and write M for shortness in the following form

$$M = X_0 y_{\sigma(1)} X_1 \cdots X_{m-1} y_{\sigma(m)} X_m$$

where X_0, \dots, X_m are some words (possibly empty) consisting of left and right brackets and indeterminates x_1, \dots, x_k . Now we define a monomial \widetilde{M} on even indeterminates x_1, \dots, x_k and odd indeterminates y_1, \dots, y_m from free Lie superalgebra as

$$\widetilde{M} = \text{sgn}(\sigma) X_0 y_{\sigma(1)} X_1 \cdots X_{m-1} y_{\sigma(m)} X_m.$$

Extending this map \sim by linearity we obtain a linear isomorphism $P_{k,m} \rightarrow P_{k,m}$ of two subspaces of a \mathbb{Z}_2 -graded free Lie algebra and a free Lie superalgebra, respectively. Although the monomials in $P_{k,m}$ are not linearly independent, it easily follows from Jacobi and super-Jacobi identities that the map \sim is well-defined. Similarly, we can define the inverse map from a free Lie superalgebra to a free \mathbb{Z}_2 -graded Lie algebra.

Following the same argument as in the associative case (see [10, Lemma 3.4.7]) we obtain for any \mathbb{Z}_2 -graded Lie algebra L and its Grassmann envelope $G(L) = G_0 \otimes L_0 \oplus G_1 \otimes L_1$ the following result.

Lemma 2. *Let $f \in P_{k,m}$ be a multilinear polynomial in the free Lie algebra. Then*

- *f is a graded identity of L if and only if \widetilde{f} is a graded identity of $G(L)$; and*
- *$\widetilde{\widetilde{f}} = f$.* □

The next lemma is an obvious generalization of Lemma 1.

Lemma 3. *Let $L = L_0 \oplus L_1$ be a finite dimensional Lie algebra, $\dim L_0 = k$, $\dim L_1 = l$, and let*

$$\chi_{q,n-q}(L) = \sum_{\substack{\lambda \vdash q \\ \mu \vdash n-q}} m_{\lambda,\mu} \chi_{\lambda,\mu}$$

be its $(q, n-q)$ -graded cocharacter. If $m_{\lambda,\mu} \neq 0$ then $\lambda \in H(k, 0)$ and $\mu \in H(l, 0)$. □

Using this remark we restrict the shape of the graded cocharacter of the Grassmann envelope $G(L)$.

Lemma 4. *Let $L = L_0 \oplus L_1$ be a finite dimensional Lie algebra, $\dim L_0 = k$, $\dim L_1 = l$, and let \tilde{L} be its Grassmann envelope. If*

$$(8) \quad \chi_{q,n-q}(\tilde{L}) = \sum_{\substack{\lambda \vdash q \\ \mu \vdash n-q}} m_{\lambda,\mu} \chi_{\lambda,\mu}$$

and $m_{\lambda,\mu} \neq 0$ in (8) then $\lambda \in H(k, 0)$ and $\mu \in H(0, l)$.

Proof. Suppose $m_{\lambda,\mu} \neq 0$ in (8) for some $\lambda \vdash q, \mu \vdash n - q$. Then there exists a multilinear polynomial $g = g(x_1, \dots, x_q, y_1, \dots, y_{n-q})$ such that

$$f = e_{T_\lambda} e_{T_\mu} g(x_1, \dots, y_{n-q})$$

is not a graded identity of \tilde{L} , where $e_{T_\lambda} \in FS_q, e_{T_\mu} \in FS_{n-q}$ are essential idempotents generating minimal left ideals in FS_q, FS_{n-q} , respectively. Inclusion $\lambda \in H(k, 0)$ immediately follows by Lemma 3 since L and $G(L)$ have the same cocharacters on even indeterminates. Since e_{T_λ} and e_{T_μ} commute, applying Lemma 4.8.6 from [10] we get

$$\tilde{f} = ae_{T_\lambda} g,$$

where $a \in I_{\mu'}$. Here μ' is the conjugated to μ partition of $n - q$ and $I_{\mu'}$ is the minimal two-sided ideal of FS_{n-q} generated $e_{T_{\mu'}}$. That is, $I_{\mu'}$ has the character $r \cdot \chi_{\mu'}$, where $r = d_{\mu'} = \deg \mu'$.

By Lemma 2, \tilde{f} is not a graded identity of $G(\tilde{L})$. Since $\tilde{h} = h$ for any $h \in P_{q,n-q}$, we see that \tilde{f} is not a graded identity of L and $\mu' \in H(l, 0)$ by Lemma 3. In other words, the number of rows of Young diagram $D_{\mu'}$ does not exceed l . This number equals the number of columns of D_μ hence $\mu \in H(0, l)$ and we are done. \square

Using the previous lemma we restrict the shape of non-graded cocharacter of $G(L)$.

Lemma 5. *Let $L = L_0 \oplus L_1$ be a finite dimensional Lie algebra, $\dim L_0 = k$, $\dim L_1 = l$, and let*

$$\chi(\tilde{L}) = \sum_{\lambda \vdash n} m_\lambda \chi_\lambda$$

be the n -th (non-graded) cocharacter of $\tilde{L} = G(L)$. Then $m_\lambda \neq 0$ only if $\lambda \in H(k, l)$.

Proof. Suppose $f \in P_n$ is not an identity of \tilde{L} . Since f is multilinear we may assume that $f(x_1, \dots, x_q, y_1, \dots, y_{n-q}) \in P_{q,n-q}$ is not an identity of \tilde{L} for some $0 \leq q \leq n$. Moreover, we can consider only the case when a graded polynomial f generates in $P_{q,n-q}$ an irreducible $S_q \times S_{n-q}$ -submodule M with the character (χ_λ, χ_μ) , $\lambda \vdash q, \mu \vdash n - q$.

Now we lift the $S_q \times S_{n-q}$ -action up to an S_n -action and consider a decomposition of $FS_n M$ into irreducible components:

$$\chi(FS_n M) = \sum_{\nu \vdash n} m_\nu \chi_\nu.$$

Since λ lies in the hook $H(k, 0)$, i.e. the horizontal strip of height k by Lemma 4 and μ lies in $H(0, l)$, the vertical strip of width l , it follows from the Littlewood-Richardson rule for induced representations ([13, 2.8.13], see also [10, Theorem 2.3.9]) that $m_\nu = 0$ as soon as $\nu \notin H(k, l)$ and we have completed the proof. \square

Lemma 6. *Let $G(L) = \tilde{L} = \tilde{L}_0 \oplus \tilde{L}_1$ be the Grassmann envelope of a finite dimensional Lie algebra $L = L_0 \oplus L_1$ with $\dim L_0 = k, \dim L_1 = l$. Then its colength sequence $\{l_n(\tilde{L})\}$ is polynomially bounded.*

Proof. We use the notation $\{z_1, z_2, \dots\}$ for non-graded indeterminates here since $\{x_1, x_2, \dots\}$ were even variables in the previous statements.

Let

$$(9) \quad \chi(\tilde{L}) = \sum_{\lambda \vdash n} m_\lambda \chi_\lambda$$

be the n -th cocharacter of \tilde{L} . By Lemma 5 we have $\lambda \in H(k, l)$ as soon as $m_\lambda \neq 0$ in (9). Fix $\lambda \vdash n$ with $m_\lambda = m \neq 0$ and consider the FS_n -submodule

$$(10) \quad W_1 \oplus \dots \oplus W_m \subseteq P_n(\tilde{L})$$

with $\chi(W_i) = \chi_\lambda$, for all $i = 1, \dots, m$.

We shall prove that

$$(11) \quad m \leq (k + l)2^{kl}n^{k^2+l^2}$$

in (10). Denote by $\lambda'_1, \dots, \lambda'_l$ the heights of the first l columns of the Young diagram D_λ . Clearly, it suffices to prove the inequality (11) only for λ with $\lambda_k > l$ and $\lambda'_l > k$. Otherwise $\lambda \in H(k', l')$ with $k' \leq k, l' \leq l$ and $k' + l' < k + l$.

Denote

$$\mu_1 = \lambda'_1 - k, \dots, \mu_l = \lambda'_l - k.$$

Then $\lambda_1 + \dots + \lambda_k + \mu_1 + \dots + \mu_l = n$.

It is well-known (see, for example, [29]) that one can choose multilinear $f_1 \in W_1, \dots, f_m \in W_m$ such that $FS_n f_1 = W_1, \dots, FS_n f_m = W_m$ and each $f_i, i = 1, \dots, m$, is symmetric on k sets of indeterminates of orders $\lambda_1, \dots, \lambda_k$ and is alternating on l sets of orders μ_1, \dots, μ_l .

According to this decomposition into symmetric and alternating sets we rename z_1, \dots, z_n as follows

$$(12) \quad \{z_1, \dots, z_n\} = \{z_1^1, \dots, z_{\lambda_1}^1, \dots, z_1^k, \dots, z_{\lambda_k}^k, \bar{z}_1^1, \dots, \bar{z}_{\mu_1}^1, \dots, \bar{z}_1^l, \dots, \bar{z}_{\mu_l}^l\},$$

where each f_i is symmetric on any set $\{z_1^j, \dots, z_{\lambda_j}^j\}, j = 1, \dots, k$, and is alternating on any set $\{\bar{z}_1^s, \dots, \bar{z}_{\mu_s}^s\}, s = 1, \dots, l$.

We shall find $\delta_1, \dots, \delta_m \in F$ such that

$$f = \delta_1 f_1 + \dots + \delta_m f_m$$

is an identity of \tilde{L} if (11) does not hold. Note that for any $\delta_1, \dots, \delta_m \in F$ a polynomial f is also symmetric on each subset $\{z_1^i, \dots, z_{\lambda_i}^i\}, 1 \leq i \leq k$, and alternating on each subset $\{\bar{z}_1^s, \dots, \bar{z}_{\mu_s}^s\}, s = 1, \dots, l$.

Let $E = \{e_1, \dots, e_{k+l}\}$ be a homogeneous basis of L with $E_0 = \{e_1, \dots, e_k\} \subset L_0, E_1 = \{e_{k+1}, \dots, e_{k+l}\} \subset L_1$. Then f is an identity of \tilde{L} if and only if $\varphi(f) = 0$ for any evaluation $\varphi: Z \rightarrow \tilde{L}$ such that $\varphi(z_i) = g_i \otimes a_i, 1 \leq i \leq n$, where a_i is a basis element from E and $g_i \in G$ has the same parity as a_i and $g_1 \cdots g_n \neq 0$ in G .

Note also that $\varphi(f) = 0$ implies $\varphi'(f) = 0$ for any evaluation φ' such that $\varphi'(z_i) = g'_i \otimes a_i, 1 \leq i \leq n$ provided that $g_1 \cdots g_n \neq 0$.

Using these two remarks we shall find an upper bound for the number of evaluations for asking the question whether f is an identity of \tilde{L} or not.

Consider first one symmetric subset $Z_1 = \{z_1^1, \dots, z_{\lambda_1}^1\}$. If $\varphi(z_i^1) = g \otimes e, \varphi(z_j^1) = h \otimes e$, for some $i \neq j$ with $e \in E_1$, then $\varphi(f) = 0$, as follows from the symmetry on Z_1 . Hence we need to check only evaluations with at most $r \leq l$ odd values $\varphi(z_{i_1}^1) = g_1 \otimes e_{t_1}, \dots, \varphi(z_{i_r}^1) = g_r \otimes e_{t_r}$, where $e_{t_1}, \dots, e_{t_r} \in E_1$ are distinct. Since Z_1 is the symmetric set of variables, the result of evaluation φ does not depend (up to the sign) on the choice of i_1, \dots, i_r . Hence we have $\binom{l}{r}$ possibilities.

Given $0 \leq r \leq l$, we estimate the number of evaluations of remaining $\lambda_1 - r$ variables in the even component of \tilde{L} . First, let $r = 0$ and $\varphi(z_i^1) = g_i \otimes a_i, a_i \in E_0, 1 \leq i \leq \lambda_1$. If e_1 appears in the row $(a_1, \dots, a_{\lambda_1})$ exactly α_1 times, e_2 appears α_2 times and so on, then the result of such substitution depends only on $\alpha_1, \dots, \alpha_k$ since f is symmetric on Z_1 . Hence we have no more than $(\lambda_1 + 1)^k$ variants since $0 \leq \alpha_1, \dots, \alpha_k \leq \lambda_1$. In particular, we need at most $(n + 1)^k$ evaluations if $r = 0$.

Let now $r = 1$. We can replace by odd element an arbitrary variable from Z_1 and get (up to the sign) the same value $\varphi(f)$ since f is symmetric on Z_1 . Suppose say, that $\varphi(z_{\lambda_1}^1) = h \otimes e, e \in E_1$, and $\varphi(z_1^1) = g_1 \otimes a_1, \dots, \varphi(z_{\lambda_1-1}^1) = g_{\lambda_1-1} \otimes a_{\lambda_1-1}$, where all a_j are even. If $\alpha_1, \dots, \alpha_k$ are the same integers as in the case $r = 0$ then the result of the substitution also depends only on $\alpha_1, \dots, \alpha_k$. Hence for $r = 1$ we have at most

$$\binom{l}{1} \lambda_1^k \leq \binom{l}{1} (n + 1)^k$$

variants for φ since $0 \leq \alpha_1, \dots, \alpha_k \leq \lambda_1 - 1$.

Similarly, for general $0 \leq r \leq l$ we have at most

$$\binom{l}{r} (\lambda_1 + 1 - r)^k \leq \binom{l}{r} (n + 1)^k$$

variants. Therefore for evaluating all variables from Z_1 it suffices

$$\sum_{r=0}^l \binom{l}{r} (n + 1)^k = 2^l (n + 1)^k$$

substitutions and for all symmetric variables we need at most

$$(2^l (n + 1)^k)^k$$

substitutions.

Now consider the alternating set $Z'_1 = \{\bar{z}_1^1, \dots, \bar{z}_{\mu_1}^1\}$. If $\varphi(\bar{z}_i^1) = g \otimes e, \varphi(\bar{z}_j^1) = h \otimes e$, for some $i \neq j$ with the same $e \in E_0$, then $\varphi(f) = 0$, hence we can choose only $0 \leq r \leq k$ distinct basis elements $b_1, \dots, b_r \in E_0$ for values of $\bar{z}_{i_1}^1, \dots, \bar{z}_{i_r}^1$ of the type $g_i \otimes b_i$. Up to the sign, the result of the substitution does not depend on i_1, \dots, i_r and we have only $\binom{k}{r}$ options.

Suppose now that all $\varphi(\bar{z}_i^1), 1 \leq i \leq r$, are fixed even values. Let

$$\varphi(\bar{z}_{r+1}^1) = g_1 \otimes b_1, \dots, \varphi(\bar{z}_{\mu_1}^1) = g_{\mu_1-r} \otimes b_{\mu_1-r}, \quad b_1, \dots, b_{\mu_1-r} \in E_1.$$

Then (up to the sign) the result of φ depends only on the number of entries of e_{k+1}, \dots, e_{k+l} into the row $(b_1, \dots, b_{\mu_1-r})$. Hence we have at most $(\mu_1 - r + 1)^l$ variants for substitution of odd variables. As in the symmetric case we have the following upper bound

$$\sum_{r=0}^k \binom{k}{r} (n + 1)^l = 2^k (n + 1)^l$$

for one subset and $(2^k (n + 1)^l)^l$ for all skew variables.

We have proved that one can find $T \leq 2^{kl}(n+1)^{l^2+k^2}$ evaluations $\varphi_1, \dots, \varphi_T$ such that the relations

$$(13) \quad \varphi_1(f) = \dots = \varphi_T(f) = 0$$

imply $\varphi(f) = 0$ for any evaluation φ , that is f is an identity of \tilde{L} . Recall that $f = \delta_1 f_1 + \dots + \delta_m f_m$. Therefore for any evaluation φ the equality $\varphi(f) = 0$ can be viewed as a system of $k+l$ homogeneous linear equations in the algebra \tilde{L} on unknown coefficients $\delta_1, \dots, \delta_m$. If (11) does not hold then the system (13) has a non-trivial solution $\bar{\delta}_1, \dots, \bar{\delta}_m$ and $f = \bar{\delta}_1 f_1 + \dots + \bar{\delta}_m f_m$ is an identity of \tilde{L} , a contradiction.

We have proved the inequality (11). From this inequality it follows that all multiplicities in (9) are bounded by $(k+l)2^{2kl}n^{k^2+l^2}$. Finally note that the number of partitions $\lambda \in H(k, l)$ is bounded by n^{k+l} . Hence

$$l_n(\tilde{L}) < (k+l)2^{2kl}n^{k^2+l^2+kl}$$

and we have thus completed the proof. \square

As a corollary of previous results we obtain the following:

Proposition 1. *Let $L = L_0 \oplus L_1$ be a finite dimensional \mathbb{Z}_2 -graded Lie algebra with $\dim L_0 = k, \dim L_1 = l$ and let $\tilde{L} = G(L)$ be its Grassmann envelope. Then there exist constants $\alpha, \beta \in \mathbb{R}$ such that*

$$c_n(\tilde{L}) \leq \alpha n^\beta (k+l)^n.$$

In particular,

$$\overline{\exp}(\tilde{L}) = \limsup_{n \rightarrow \infty} \sqrt[n]{c_n(\tilde{L})} \leq k+l.$$

Proof. By [10, Lemma 6.2.5], there exist constants C and r such that

$$\sum_{\lambda \in H(k, l)} d_\lambda \leq C n^r (k+l)^n$$

for all $n = 1, 2, \dots$. In particular,

$$\max\{d_\lambda | \lambda \vdash n, \lambda \in H(k, l)\} \leq C n^r (k+l)^n.$$

Now Lemma 6 and the inequality (6) complete the proof. \square

4. EXISTENCE OF PI-EXPONENTS

Proposition 2. *Let L be a finite dimensional simple Lie algebra over an algebraically closed field of characteristic zero with some \mathbb{Z}_2 -grading, $L = L_0 \oplus L_1$, $\dim L_0 = k, \dim L_1 = l$. Let also $\tilde{L} = G(L)$ be its Grassmann envelope. Then there exist constants $\gamma > 0, \delta \in \mathbb{R}$ such that*

$$c_n(\tilde{L}) \geq \gamma n^\delta (k+l)^n.$$

In particular,

$$\underline{\exp}(\tilde{L}) = \liminf_{n \rightarrow \infty} \sqrt[n]{c_n(\tilde{L})} \geq k+l.$$

Proof. Denote $d = k + l = \dim L$. By [19, Theorem 12.1], for the adjoint representation of L there exists a multilinear associative polynomial $h = h(u_1^1, \dots, u_d^1, \dots, u_1^m, \dots, u_d^m)$ alternating on each subset of indeterminates $\{u_1^i, \dots, u_d^i\}$, $1 \leq i \leq m$, such that under any evaluation $\varphi : u_j^i \rightarrow ad \ b_j^i, b_j^i \in L$, the value $\varphi(h)$ is a scalar linear transformation of L and $\varphi(h) \neq 0$ for some h . It follows that for any integer $t \geq 1$ there exists a multilinear Lie polynomial

$$f_t = f_t(u_1^1, \dots, u_d^1, \dots, u_1^{mt}, \dots, u_d^{mt}, w)$$

alternating on each set $\{u_1^i, \dots, u_d^i\}$, $1 \leq i \leq mt$ such that $\varphi(f_t) \neq 0$ for some evaluation $\varphi : \{u_1^1, \dots, u_d^{mt}, w\} \rightarrow L_0 \cup L_1$. Since f_t is multilinear and alternating on each set $\{u_1^i, \dots, u_d^i\}$ and $d = \dim L_0 + \dim L_1$ it follows that for any $t \geq 1$ we get a graded multilinear polynomial

$$f_t = f_t(x_1^1, \dots, x_k^1, \dots, x_1^{mt}, \dots, x_k^{mt}, y_1^1, \dots, y_l^1, \dots, y_1^{mt}, \dots, y_l^{mt}, w)$$

which is not a graded identity of L and it is alternating on each subset $\{x_1^i, \dots, x_k^i\}$ and on each subset $\{y_1^i, \dots, y_l^i\}$, $1 \leq i \leq mt$, where x_j^i 's are even and y_j^i 's are odd variables. The latter indeterminate w can be taken of arbitrary parity, say, $w = x_0$ is even.

Consider an $S_p \times S_q$ -action on

$$P_{p+1,q} = P_{p+1,q}(x_0, x_1^1, \dots, x_k^{mt}, y_1^1, \dots, y_l^{mt}),$$

where $p = mtk, q = mtl$ and S_p, S_q act on $\{x_j^i\}, \{y_j^i\}$, respectively. It follows from Lemma 3 that the $S_p \times S_q$ -character of the submodule generated by f in $P_{p+1,q}$ lies in the pair of of strips $H(k, 0), H(l, 0)$, that is

$$\chi(F[S_p \times F_q]f) = \sum_{\substack{\lambda \vdash p \\ \mu \vdash q}} m_{\lambda,\mu} \chi_{\lambda,\mu}$$

with $m_{\lambda,\mu} = 0$, unless $\lambda \in H(k, 0), \mu \in H(l, 0)$. Hence λ is a partition of mtk with at most k rows. On the other hand, f depends on mt alternating subsets of even indeterminates of order k each. It is well-known that in this case $m_{\lambda,\mu} = 0$ if $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\lambda_1 \geq mt + 1$. It follows that only rectangular partition

$$(14) \quad \lambda = (\underbrace{mt, \dots, mt}_k)$$

can appear in $F[S_p \times F_q]f$ with non-zero multiplicity. Similarly,

$$(15) \quad \mu = (\underbrace{mt, \dots, mt}_l)$$

if $m_{\lambda,\mu} \neq 0$. Hence we can assume that f has the form

$$f = e_{T_\lambda} e_{T_\mu} g(x_1^1, \dots, y_l^{mt}, w)$$

with λ and μ of the types (14), (15), respectively.

By Lemma 2, the polynomial \tilde{f} is not an identity of the Lie superalgebra $\tilde{L} = G(L)$ and by Lemma 4.8.6 from [10], the graded polynomial \tilde{f} generates in $P_{p+1,q}(\tilde{L})$ an irreducible $S_p \times S_q$ -submodule with the character $(\chi_\lambda, \chi_{\mu'})$, where

$$\mu' = (\underbrace{l, \dots, l}_{mt})$$

is conjugated to a μ partition of mtl .

First we apply Littelwood-Richardson rule and induce this $S_p \times S_q$ -module up to S_n -module. Then we induce the obtained S_n -module up to S_{n+1} -module, where $n = p + q = mt(k + l)$. It follows from the Littelwood-Richardson rule that the induced S_{n+1} -module can contain only simple submodule corresponding to partitions $\nu \vdash n + 1$ such that the Young diagram D_ν contains a subdiagram D_{ν_0} , where

$$\nu_0 = h(k, l, t_0) = (\underbrace{l + t_0, \dots, l + t_0}_k, \underbrace{l, \dots, l}_{t_0})$$

is a finite hook with $t_0 \geq l - k, mt - kl$. Since we are interested with asymptotic of codimensions, we may assume that $mt - kl > l - k$ and then $t_0 = mt - kl$. In particular, ν_0 is a partition of $n_0 = (k + l)t_0 + kl$. Then $n + 1 - n_0 = (k + l - 1)kl + 1$ and by [10, Lemma 6.2.4]

$$d_{\nu_0} \leq d_\nu \leq n^c d_{\nu_0}$$

where $c = (k + l - 1)kl + 1$ and

$$d_{h(k, l, t_0)} \simeq an_0^b (k + l)^{n_0} \quad \text{if } n_0 \rightarrow \infty$$

for some constants a, b , by Lemma 6.2.5 from [10]. Here the relation $f(n) \simeq g(n)$ means that $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 1$. Since $c_{n+1}(\tilde{L}) \geq d_\nu$ we get the inequality

$$(16) \quad c_{n+1}(\tilde{L}) \geq \alpha(n + 1)^\beta (k + l)^{n+1}$$

for all $n = m(k + l)t$, $t = 1, 2, \dots$ for some constants $\alpha > 0$ and β .

Since Lie algebra L is simple, the Grassmann envelope \tilde{L} is a centerless Lie superalgebra. It is not difficult to see that $c_{r+1}(\tilde{L}) \geq c_r(\tilde{L})$ in this case for all $r \geq 1$. Hence by (16) we have

$$c_{n+j}(\tilde{L}) \geq \alpha(n + 1)^\beta (k + l)^{n+1}$$

for any $1 \leq j \leq m(k + l)$. Since $n = m(k + l)t$ one can find constants $\gamma > 0$ and δ such that

$$c_r(\tilde{L}) \geq \gamma r^\delta (k + l)^r$$

for all positive integer r and we have completed the proof. \square

Theorem 1 now easily follows from Propositions 1 and 2.

Proof of Theorem 2. First we obtain an upper bound for $c_n^{gr}(\tilde{L})$:

$$c_n^{gr}(\tilde{L}) = \sum_{q=0}^n \binom{n}{q} c_{q, n-q}(\tilde{L}),$$

where

$$(17) \quad c_{q, n-q}(\tilde{L}) = \sum_{\substack{\lambda \vdash q \\ \mu \vdash n-q}} m_{\lambda, \mu} d_{\lambda, \mu}$$

and $d_{\lambda, \mu} = \deg \chi_{\lambda, \mu} = \deg \chi_\lambda \cdot \deg \chi_\mu = d_\lambda d_\mu$. Moreover, $\lambda \in H(k, 0)$, $\mu \in H(0, l)$ by Lemma 4. Applying Lemma 6.2.5 from [10], we obtain

$$\sum_{\substack{\lambda \in H(k, 0) \\ \lambda \vdash q}} d_\lambda \leq C n^r k^q, \quad \sum_{\substack{\mu \in H(0, l) \\ \mu \vdash n-q}} d_\mu \leq C n^r l^{n-q}$$

for some constants C, r and hence

$$(18) \quad \sum_{\substack{\lambda \in H(k, 0), \lambda \vdash q \\ \mu \in H(0, l), \mu \vdash n-q}} d_\lambda d_\mu \leq C^2 n^{2r} k^q l^{n-q}$$

On the other hand, graded colength

$$l_{q,n-q}(\tilde{L}) = \sum_{\substack{\lambda \vdash q \\ \mu \vdash n-q}} m_{\lambda,\mu}$$

is not greater than non-graded colength $l_n(\tilde{L})$. Since $l_n(\tilde{L})$ is polynomially bounded by Lemma 6, one can find a polynomial $\varphi(n)$ such that

$$(19) \quad m_{\lambda,\mu} \leq \varphi(n)$$

for any $m_{\lambda,\mu}$ in (17). It now follows from (17), (18) and (19) that for $\psi(n) = C^2 n^{2r} \varphi(n)$ we have

$$(20) \quad c_n^{gr}(\tilde{L}) \leq \psi(n) \sum_{q=1}^n \binom{n}{q} k^q l^{n-q} = \psi(n)(k+l)^n$$

and we have obtained an upper bound for $c_n^{gr}(\tilde{L})$.

On the other hand, in [2, Lemma 3.1] it is proved that for any associative G -graded algebra A , where G is a finite group, an ordinary n -th codimension is less than or equal to the graded n -th codimension, for any n . Proof of this lemma does not use associativity. Hence

$$(21) \quad c_n^{gr}(\tilde{L}) \geq c_n(\tilde{L}).$$

Theorem 2 now follows from (20), (21) and Proposition 2 and we have completed the proof. \square

REFERENCES

- [1] Yu. A. Bahturin and V. Drensky. Graded polynomial identities of matrices. *Linear Algebra Appl.*, 357:15–34, 2002.
- [2] Yu. Bahturin, A. Giambruno and D. Riley. Group-graded algebras with polynomial identity. *Israel J. Math.*, 104:145–155, 1998.
- [3] A. Berele and A. Regev. Applications of hook Young diagrams to P.I. algebras. *J. Algebra*, 82(2):559–567, 1983.
- [4] A. Berele and A. Regev. On the codimensions of the verbally prime P.I. algebras. *Israel J. Math.*, 91:239–247, 1995.
- [5] A. Giambruno, S. P. Mishchenko and M. V. Zaicev. Codimensions of algebras and growth functions. *Adv. Math.*, 217(3):1027–1052, 2008.
- [6] A. Giambruno, A. Regev and M. V. Zaicev. On the codimension growth of finite-dimensional Lie algebras. *J. Algebra*, 220(2):466–474, 1999.
- [7] A. Giambruno, A. Regev and M. V. Zaicev. Simple and semisimple Lie algebras and codimension growth. *Trans. Amer. Math. Soc.*, 352(4):1935–1946, 2000.
- [8] A. Giambruno and M. Zaicev. On codimension growth of finitely generated associative algebras. *Adv. Math.*, 140:145–155, 1998.
- [9] A. Giambruno and M. Zaicev. Exponential codimension growth of P.I. algebras: an exact estimate. *Adv. Math.*, 142:221–243, 1999.
- [10] A. Giambruno and M. Zaicev. Polynomial Identities and Asymptotic Methods, Mathematical Surveys and Monographs, 122. American Mathematical Society, Providence, RI, 2005.
- [11] A. Giambruno and M. Zaicev. Codimension growth of special simple Jordan algebras. *Trans. Amer. Math. Soc.*, 362:3107–3123, 2010.
- [12] A. Giambruno and M. Zaicev. On codimension growth of finite dimensional Lie superalgebras. *J. Lond. Math. Soc.*, 85:534–548, 2012.
- [13] G. James and A. Kerber. The Representation Theory of the Symmetric Group. Encyclopedia of Mathematics and its Applications, 16. Addison-Wesley, London, 1981.
- [14] A. R. Kemer. Ideals of Identities of Associative Algebras. Translations of Mathematical Monographs, 87. American Mathematical Society, Providence, RI, 1991.

- [15] S. P. Mishchenko. On the problem of the Engel property. (Russian). *Mat. Sb.*, 124:56–67, 1984.
- [16] S. P. Mishchenko. Growth of varieties of Lie algebras. (Russian). *Uspekhi Mat. Nauk*, 45:25–45, 1990; English translation in *Russian Math. Surveys*, 45:27–52, 1990.
- [17] S. P. Mishchenko and P. M. Petrogradsky. Exponents of varieties of Lie algebras with a nilpotent commutator subalgebra. *Comm. Alg.*, 27:2223–2230, 1999.
- [18] V. M. Petrogradsky. Scale for codimension growth of Lie algebras. In *em Methods in Lie Theory* (Levico Terme, 1997), pages 213–222. Lecture Notes in Pure and Appl. Math., 198. Marcel Dekker, New York, 1998.
- [19] Yu. P. Razmyslov. Identities of algebras and their representations. Translations of Mathematical Monographs, 138. American Mathematical Society, Providence, RI, 1994.
- [20] A. Regev. Existence of identities in $A \otimes B$. *Israel J. Math.*, 11:131–152, 1972.
- [21] A. Regev. Codimensions and thace codimensions of matrices are asymptotically equal. *Israel J. Math.*, 47:246–250, 1984.
- [22] Ya. Vais. Special varieties of Lie algebras. (Russian). *Algebra i Logika*, 28(1):29–40, 1989; English translation in *Algebra and Logic*, 28(1):17–24, 1989.
- [23] M. Zaicev. Identities of affine Kac-Moody algebras. (Russian). *Vestnik Moscov. Univ. Ser. I Mat. Mekh.*, 2:33–36, 1996; English translation in *Moscow Univ Math. Bull.*, 51:29–31, 1996.
- [24] M. Zaicev. Varieties of affine Kac-Moody algebras. (Russian). *Matem. Zametki*, 62:95–102, 1997.
- [25] M. Zaicev. Integrality of exponents of growth of identities of finite-dimensional Lie algebras. (Russian). *Izv. Ross. Akad. Nauk Ser. Mat.*, 66:23–48, 2002; English translation in *Izv. Math.*, 66:463–487, 2002.
- [26] M. V. Zaitsev and S. P. Mishchenko. Varieties of Lie superalgebras of polynomial growth. (Russian). *Uspekhi Mat. Nauk*, 52(2):165–166, 1997; English translation in *Russian Math. Surveys*, 52(2):432–433, 1997.
- [27] M. V. Zaitsev and S. P. Mishchenko. A criterion for the polynomial growth of varieties of Lie superalgebras. (Russian). *Izv. Ross. Akad. Nauk Ser. Mat.*, 62(5):103–116, 1998; English translation in *Izv. Math.*, 62(5):953–967, 1998.
- [28] M. V. Zaicev and S. P. Mishchenko. An example of a variety of Lie algebras with a fractional exponent. *Journal of Mathematical Sciences*, 93:977–982, 1999.
- [29] M. V. Zaitsev and S. P. Mishchenko. On the colength of varieties of linear algebras. (Russian). *Mat. Zametki*, 79(4):3–18, 2007; English translation in *Math. Notes*, 79(3–4):511–517, 2006.
- [30] M. V. Zaitsev and S. P. Mishchenko. The growth of some varieties of Lie superalgebras. (Russian). *Izv. Ross. Akad. Nauk Ser. Mat.*, 71(4):3–18, 2007; English translation in *Izv. Math.*, 71(4):657–672, 2007.
- [31] M. V. Zaitsev and S. P. Mishchenko. Identities of Lie superalgebras with a nilpotent commutator. (Russian). *Algebra i Logika*, 47(5):617–645, 2008; English translation in *Algebra Logic* 47(5), 2008.

DUŠAN REPOVŠ, FACULTY OF MATHEMATICS AND PHYSICS, AND FACULTY OF EDUCATION, UNIVERSITY OF LJUBLJANA, P. O. B. 2964, LJUBLJANA, 1001, SLOVENIA

E-mail address: dusan.repovs@guest.arnes.si

MIKHAIL ZAICEV, DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND MECHANICS, MOSCOW STATE UNIVERSITY, MOSCOW, 119992 RUSSIA

E-mail address: zaicevmv@mail.ru